

2008-2009

**WINTER STEELHEAD
HARVEST MANAGEMENT PLAN
for the
PORT ANGELES and SEQUIM – PORT TOWNSEND
MANAGEMENT UNITS
of the
EASTERN STRAIT OF JUAN DE FUCA**

DRAFT

Joint Report Prepared by:

**Washington Department of Fish and Wildlife
Point No Point Treaty Council
(Jamestown S’Klallam Tribe - Port Gamble S’Klallam Tribe)
Lower Elwha Klallam Tribe**

October 2008

INTRODUCTION

The Washington Department of Fish and Wildlife (WDFW), the Point No Point Treaty Council (on behalf of the Jamestown S'Klallam Tribe and the Port Gamble S'Klallam Tribe), and the Lower Elwha Klallam Tribe agree on the following harvest management plan for the 2008-09 winter fishing seasons in the eastern Strait of Juan de Fuca. This plan outlines the management guidelines for the steelhead resources of two management units comprising several small independent streams between the Elwha River and Admiralty Inlet.

PREMISES AND FISHERY OBJECTIVES

The provisions of this plan cover all treaty and non-treaty fisheries for steelhead occurring in the Port Angeles and the Sequim – Port Townsend MU streams and nearshore marine areas. The co-managers agree to a philosophy of cooperation in implementing management programs to maintain, perpetuate and enhance the steelhead resource and the natural ecosystem that supports it.

This plan is intended to ensure that treaty and non-treaty fishers, subject to their respective regulatory authorities, shall be afforded the opportunities to harvest their shares as determined in *United States v. Washington*, 384 F. Supp. 312, aff'd 520 F.2d 676 (9th Cir. 1975), cert. denied 423 U.S. 1086, aff'd sub nom *Washington v. Washington State Commercial Passenger Fishing Vessel Association*, 443 U.S. 658 (1979) and other orders under the court's continuing jurisdiction.

The co-managers agree to enact and recommend for enactment by the Pacific Fishery Management Council and the Pacific Salmon Commission, appropriate regulations for marine salmonid fisheries that will provide for adequate escapement of steelhead into the Eastern Strait of Juan de Fuca watersheds, to achieve the goals and purposes of this plan.

MANAGEMENT UNITS

This Plan covers two Management Units (MUs) in the Eastern Strait of Juan de Fuca, each composed of a number of sub populations. The MUs, and their components, along with available information, are described as follows:

Port Angeles Management Unit:

The Port Angeles MU includes the aggregate of sub populations in independent streams between the Elwha River and the Dungeness River. Most sub populations are presumed to be composed of wild steelhead aggregations. The sub populations in this unit are;

Morse Creek; Available information on this sub population is outlined in Table 1

Siebert Creek Available information is outlined in Table 2

McDonald Creek Available information is outlined in Table 2.

Table 1. Total Winter Steelhead Harvest, Wild Escapement Index and Hatchery – Reared Smolt Releases in Morse Creek.

Winter Return Year	Recreational Harvest a/	Treaty Harvest	Total Harvest	Index Escapement Estimate	Smolt Year	Smolts Stocked
1980-81	357	166	523		1979	12,826
1981-82	290	116	406		1980	12,325
1982-83	310	258	568		1981	18,014
1983-84	546	216	762		1982	15,420
1984-85	285	351	636	145	1983	16,388
1985-86	235	281	516	105	1984	15,460
1986-87	223	200	423	118	1985	15,852
1987-88	188	283	471	138	1986	18,819
1988-89	215	58	273	60	1987	15,227
1989-90	189	45	234	78	1988	15,034
1990-91	90	0	90	91	1989	15,514
1991-92	135	0	135	100	1990	10,034
1992-93	96	12	108		1991	14,655
1993-94	195	0	195		1992	10,058
1994-95b/	117	2	119	128	1993	15,438
1995-96	300	11	311	89	1994	15,338
1996-97	150	0	150	183	1995	15,029
1997-98	73	0	73	102	1996	5,076
1998-99	53	0	53	81	1997	5,000
1999-00	110	0	110	162	1998	5,000
2000-01	49	0	49	99	1999	5,010
2001-02	145	0	145	71	2000	5,000
2002-03	33	0	33	84	2001	5,000
2003-04	36	0	36	121	2002	5,000
2004-05	166	3	169		2003	5,000
2005-06	51	3	54	124	2004	5,000
2006-07	33	0	33	118	2005	0
2007-08	10	0	10		2006	0
10 year mean 1998-99 to 2007-08			69	108		

a/ Hatchery and wild steelhead recreational harvest numbers available since 1986-87 from WDFW Catch Record Cards.

b/ Wild steelhead release regulations in recreational fishery since 1994-95.

Table 2. Wild Winter Steelhead index escapement estimates and smolt production in streams of the Port Angeles Winter Steelhead MU.

Spawning Year	McDonald Creek Index Escapement	Smolt Emigration Year	McDonald Creek smolts	Siebert Creek smolts
1985		1985		
1986		1986		
1987		1987		
1988		1988		
1989		1989		
1990		1990		
1991		1991		
1992		1992		
1993		1993		
1994		1994		
1995		1995		
1996		1996		
1997		1997		
1998	308	1998		1,626
1999	217	1999	1,861	551
2000	251	2000	3,343	937
2001	143	2001		
2002	125	2002		513
2003	63	2003		758
2004	29	2004		975
2005	89	2005		769
2006	206	2006	1,125	235
2007	63	2007	1,424	558
2008		2008	1,170	793

Sequim – Port Townsend Management Unit

The Sequim – Port Townsend MU includes the aggregate of sub- populations in independent streams between the Dungeness River and Admiralty Inlet. In this MU, streams empty into distinct embayments (Sequim Bay, Discovery Bay, Port Townsend Bay) and all associated basins lie in the “rain shadow” of the Olympic mountains. All of the sub- populations are presumed to be composed of wild steelhead aggregations. The sub- populations in this unit are;

- Jimmycomelately (JCL) Creek (Sequim Bay)
- Salmon Creek (Discovery Bay)
- Snow Creek (Discovery Bay)
- Chimacum Creek (Port Townsend Bay)

Available information is summarized in Tables 3 – 5.

WILD STEELHEAD ESCAPEMENT and SMOLT PRODUCTION

In the Port Angeles Management Unit, estimates of natural spawners are limited to index counts, in Morse Creek and McDonald Creek and more recently, in Siebert Creek. (Table 2). It is anticipated that these index enumerations will continue; however, spawner surveys have been limited to periods after March 1 in most seasons and stream flows may often be high during the recommended spawner survey period and the amount of information collected will depend on suitable river conditions. Smolt production from these spawners is largely unavailable, so direct estimates of productivity and productive capacities have not been possible. For this reason, no specific escapement goals have been established for these MUs.

The best life history information available on wild steelhead in this region comes from the Snow Creek trap (Sequim – Port Townsend MU) near river mile 0.8. The numbers of wild steelhead adults and smolts including smolt-to-adult survival rates have been documented since the late 1970s (Table 3). The numbers of wild Snow Creek steelhead adults by age and percent age composition are summarized in Tables 4 and 5, respectively. Figure 1 shows the percentage of wild adult steelhead returning to the Snow Creek trap by month from 1978 to 1994

Table 3. Number of JCL Creek and Snow Creek wild steelhead smolts, Snow Creek wild steelhead adult returns, and smolt-to-adult return rates.

Smolt Emigration Year(n)	Jimmycome lately Creek	Snow Creek							
	Total Wild Smolts	Wild Smolts Above Weir (a)	Total Wild Smolts (b)	1-salt Adults (n+1)	2-salt Adults (n+2)	3-salt Adults (n+3)	Total Wild Adults	Adults / Smolt (d)	Adults / Smolt (e)
1978		1,403	1,551	2	88	11	101	0.065	0.072
1979		892	1,040	9	98	4	111	0.107	0.124
1980		1,357	1,505	3	78	4	85	0.056	0.063
1981		1,541	1,689	0	33	4	37	0.022	0.024
1982		1,733	1,881	4	102	8	114	0.061	0.066
1983		1,270	1,418	10	130	9	149	0.105	0.117
1984		1,107	1,255	10	45	5	60	0.048	0.054
1985		2,074	2,222	5	59	18	82	0.037	0.040
1986		517	665	3	38	6	47	0.071	0.091
1987		1,861	2,009	0	21	3	24	0.012	0.013
1988		541	689	1	6	2	9	0.013	0.017
1989		1,713	1,861	0	30	4	34	0.018	0.020
1990		1,336	1,484	0	44	5	49	0.033	0.037
1991		1,162	1,310	0	21	6	27	0.021	0.023
1992		2,079	2,227	2	35	0	37	0.017	0.018
1993		1,513	1,661	0	45	8	53	0.032	0.035
1994		1,583	1,731	0	131	5	136	0.079	0.086
1995		297	445	0	52	3	55	0.124	0.185
1996		2,015	2,163	2	56	6	64	0.030	0.032
1997		1,164	1,312	0	43	4	47	0.036	0.040
1998		778	926	4	152	8	164	0.177	0.211
1999		(f)	1,478	3	44	5	52	0.035	0.039
2000		1,285	1,433	2	23	4	29	0.020	0.023
2001		2,346	2,494	0	78	5	83	0.033	0.035
2002	62	2,298	2,446	2	30	1	33	0.013	0.014
2003	105	2,589	2,737	0	14	0	14	0.005	0.005
2004	191	525	673	0	22	0	22	0.033	0.042
2005	130	1,103	1,251	?	?				
2006	91	660	808	?					
2007	133	920	1,068						
2008	108								

a/ Number of steelhead smolts collected at WDFW trap at RM 0.0

b/ Includes estimated number of steelhead smolts produced downstream of WDFW trap

c/ Number of adults includes fish trapped at trap and estimated number of adults spawning downstream of trap; includes repeat spawners

d/ Includes number of steelhead smolts collected at trap RM 0.8 plus estimate of smolts downstream of trap

e/ Includes only number of steelhead smolts collected at trap RM 0.8.

f/ No smolt count due to damaged downstream trap. For calculations, used average from 1978 to 1998 for the purpose of estimating smolt production.

Table 4. Number of wild steelhead adults in Snow Creek for the return years 1976-77 through 2006-07.

Run Year a/	1-salt b/ (x.+)	2-salt b/ (x.1+)	3-salt b/ (x.2+)	Repeat spawners b/ x.1+S+ ex.	Total Wild Escapement
1976-77	6	78	2	20	106
1977-78	2	117	2	19	140
1978-79	2	61	6	12	81
1979-80	9	88	10	12	119
1980-81	3	98	11	16	128
1981-82	0	78	4	27	109
1982-83	4	33	4	11	52
1983-84	10	102	4	15	131
1984-85	10	130	8	6	154
1985-86	5	45	9	2	61
1986-87	3	59	5	5	72
1987-88	0	38	18	15	71
1988-89	1	321	6	1	29
1989-90	0	6	3	3	12
1990-91	0	30	2	2	34
1991-92	0	44	4	3	51
1992-93	2	21	5	2	30
1993-94	0	35	6	0	41
1994-95	0	45	0	0	45
1995-96	0	131	8	0	139
1996-97	2	52	5	14	73
1997-98	0	56	3	5	64
1998-99	4	43	6	2	55
1999-00	3	152	4	7	166
2000-01	2	44	8	4	58
2001-02	0	23	5	0	28
2002-03	2	78	4	6	90
2003-04	0	30	5	5	40
2004-05	0	14	1	0	15
2005-06	0	22	0	0	22
2006-07	?	?	?	?	34


a/ Run year equals winter steelhead that return to streams between November and December of year "n" and January through May of year "n+1")

b/ The letter "x" refers to the freshwater age of steelhead at Snow Creek. The time spent in freshwater prior to the first saltwater entry is shown by an Arabic numeral and/or a "+" followed by a period. The numbers of completed winters at sea prior to the last freshwater entry (a spawning run for example) is designated by (an) Arabic numerals (s) and are placed after the period. A "completed winter" on a scale is a series of tightened circuli (annulus) followed by more widely spaced circuli (spring or summer growth). Partial years at sea are denoted by a "+". An "S" is used to designate a spawning check (scale resorption).

Table 5. Percent age composition of Snow Creek wild steelhead adults referred to in Table 4.

Run Year	Percentage of First Time and Repeat Spawners				Age Composition of First Time Spawners		
	1-salt(x.+)	2-salt x.1+	3-salt x.2+	Repeat spawners	1-salt x.+	2-salt x.1+	3-salt x.2+
1976-77	5.7%	73.6%	1.9%	18.9%	7.0%	90.7%	2.3%
1977-78	1.4%	83.6%	1.4%	13.6%	1.7%	96.7%	1.7%
1978-79	2.5%	75.3%	7.4%	14.8%	2.9%	88.4%	8.7%
1979-80	7.6%	73.9%	8.4%	10.1%	8.4%	82.2%	9.3%
1980-81	2.3%	76.6%	8.6%	12.5%	2.7%	87.5%	9.8%
1981-82	0.0%	71.6%	3.7%	24.8%	0.0%	95.1%	4.9%
1982-83	7.7%	63.5%	7.7%	21.2%	9.8%	80.5%	9.8%
1983-84	7.6%	77.9%	3.1%	11.5%	8.6%	87.9%	3.4%
1984-85	6.5%	84.4%	5.2%	3.9%	6.8%	87.8%	5.4%
1985-86	8.2%	73.8%	14.8%	3.3%	8.5%	76.3%	15.3%
1986-87	4.2%	81.9%	6.9%	6.9%	4.5%	88.1%	7.5%
1987-88	0.0%	53.5%	25.4%	21.1%	0.0%	67.9%	32.1%
1988-89	3.4%	72.4%	20.7%	3.4%	3.6%	75.0%	21.4%
1989-90	0.0%	50.0%	25.0%	25.0%	0.0%	66.7%	33.3%
1990-91	0.0%	88.2%	5.9%	5.9%	0.0%	93.8%	6.3%
1991-92	0.0%	86.3%	7.8%	5.9%	0.0%	91.7%	8.3%
1992-93	6.7%	70.0%	16.7%	6.7%	7.1%	75.0%	17.9%
1993-94	0.0%	85.4%	14.6%	0.0%	0.0%	85.4%	14.6%
1994-95	0.0%	100.0%	0.0%	0.0%	0.0%	100.0%	0.0%
1995-96	0.0%	94.2%	5.8%	0.0%	0.0%	94.2%	5.8%
1996-97	2.7%	71.2%	6.8%	19.2%	3.4%	98.1%	8.5%
1997-98	0.0%	87.5%	4.7%	7.8%	0.0%	94.9%	5.1%
1998-99	7.3%	78.2%	10.9%	3.6%	7.5%	81.1%	11.3%
1999-00	1.8%	91.6%	2.4%	4.2%	1.9%	95.6%	2.5%
2000-01	3.4%	75.9%	13.8%	6.9%	3.7%	81.5%	14.8%
2001-02	0.0%	82.1%	17.9%	0.0%	0.0%	82.1%	17.9%
2002-03	2.2%	86.7%	4.4%	6.7%	2.4%	92.9%	4.8%
2003-04	0.0%	75.0%	12.5%	12.5%	0.0%	85.7%	14.3%
2004-05	0.0%	93.3%	6.7%	0.0%	0.0%	93.3%	6.7%
2005-06	0.0%	100.0%	0.0%	0.0%	0.0%	100.0%	0.0%
2006-07							
Mean	2.7%	79.3%	9.0%	9.0%	3.0%	86.9%	10.1%
Min.	0.0%	50.0%	0.0%	0.0%	0.0%	66.7%	0.0%
Max.	8.2%	100.0%	25.4%	25.0%	9.8%	100.0%	33.3%

Figure 1. Percentage of wild adult steelhead returning to the Snow Creek trap by month from 1978 to 1994.



HATCHERY CONTRIBUTIONS

In Morse Creek hatchery reared steelhead are derived from the early timed Chambers Creek stock, and for many years was transferred from the Bogachiel Hatchery. From 1977 to 1996 approximately 15,000 hatchery steelhead smolts were planted annually into Morse Creek. To reduce the impact of hatchery steelhead on to wild steelhead stocks the hatchery steelhead smolt releases were reduced to 5,000 annually and were discontinued in 2005. All hatchery-reared steelhead are marked with an adipose fin clip to enable mark selective fishing opportunities, and monitoring of stray rates should funding become available.

MANAGEMENT AND ACCOUNTING PERIODS

The management periods indicated in this report define the time interval during which regulatory actions are directed to meeting conservation and allocation needs of steelhead runs while taking into account harvest (actual or expected) of steelhead made outside the management periods. Since many runs extend over lengthy periods of time and only a small portion of the population of each run is available at the extremes of its run timing, it is impractical to exercise directed management for non-selective fisheries on these portions of runs while continuing harvests of other species or stocks. However, effort should be spread throughout the management periods to achieve escapement and harvest from all segments of the runs.

2008 – 2009 Winter Steelhead Management Period (Both Management Units):
November 30, 2008 - April 15, 2009

Management periods for winter steelhead were established by reviewing historical information on

the timing of recreational catches. The end of the management period was adjusted to match the beginning of the spring Chinook management period.

Harvest accounting periods for all treaty Indian and recreational fisheries for winter steelhead are November 1 to April 30.

HARVEST MANAGEMENT OUTLINE FOR 2008 - 2009

For winter steelhead, there is uncertainty regarding the status of some wild stocks and the degree of run timing overlap between hatchery and naturally reared steelhead in tributaries of the Strait of Juan de Fuca. Limited available information indicates that catches after the end of February may be comprised primarily of naturally reared steelhead. In order to provide additional protection for that portion of the run, fishery related mortalities, after the end of February are being severely limited. This management strategy is expected to provide for the harvest of hatchery – reared steelhead while limiting impacts to later, likely mostly wild, steelhead entering the streams after March 1.

The great majority of streams in each of these two management units are closed to fishing, during the winter steelhead accounting period. More specifically, for the 2008 – 2009 winter steelhead accounting period, co-managers have adopted the following regulatory framework.

Port Angeles Management Unit

Morse Creek	Treaty	Open for hook and line subsistence fishing from November 30 through March 14, 2 fish bag. Commercial hook and line open from November 30, through February 28 with four fish bag. Gillnet use, when authorized, may be no more than four days per week.
	Nontreaty	Open to recreational harvest from December 1, through February 28min, size 14", 2 fish bag.
Siebert Creek	All	Closed to fishing.
McDonald Creek	All	Closed to fishing

Anticipated Total Harvest: 69 ± 53

Sequim – Port Townsend Management Unit

Jimmycomelately Creek	All	Closed to fishing
Salmon Creek	All	Closed to fishing
Snow Creek	All	Closed to fishing
Chimacum Creek	All	Closed to fishing

Anticipated Total Harvest: Zero

Net fishery openings, where appropriate, shall be announced by emergency regulations based on in-season management considerations concerning the status of the stocks.

CATCH ACCOUNTING

Harvest accounting will include all commercial (including take – home), recreational and subsistence harvest in each Management Unit. Direct harvest will be reported as appropriate, on fish receiving tickets, recreational catch record cards, and treaty subsistence record cards. The co-managers will further investigate viable methods to estimate unrecorded catches or other bycatch mortalities, not reported by the above methods.

ADDITIONAL CONSERVATION MEASURES

The Co-Managers will continue to work together towards restoring and maintaining the abundance, distribution, diversity, and long-term productivity of wild steelhead and their habitats to assure healthy, self-sustaining stocks.

- Natural Production- continue protecting natural production through recreational wild steelhead release regulations in freshwater and marine areas, close rivers to recreational steelhead fishing by March 1. Harvest of the naturally produced steelhead in the Morse Creek system is minimal and should be having little negative influence on the health and abundance of the natural population.
- Habitat Protection and Restoration- continue working with local government authorities, salmon coalitions, and the public with on-going stream habitat protection and estuarine restoration efforts on Chimacum, Salmon, and JimmyComeLately creeks. The current summer chum and coho recovery efforts in this area will enhance biodiversity and nutrient enrichment for all species, including steelhead, in these watersheds.
- Fishery Management- Recreational harvest is managed to focus on hatchery steelhead while minimizing any effect on wild steelhead and treaty harvest has ranged from 0 to 12, or less than 2 fish per season since the 1990-91 season in Morse Creek.
- Artificial Production- releases of hatchery steelhead have occurred only in the Morse Creek watershed. Hatchery steelhead smolt releases in Morse Creek averaged 15,000 from 1977 to 1994 but decreased to 5,000 annually to reduce impacts to wild fish. Hatchery reared steelhead releases were discontinued in 2006.
- Monitoring, Evaluation, and Adaptive Management-WDFW will continue to monitor wild steelhead adult and smolt numbers at the Snow Creek weir and conduct steelhead spawner surveys in Morse Creek and McDonald Creek. The following creeks are currently being monitored for salmonid smolt production by the Jamestown S'Klallam Tribe:
JimmyComeLately, McDonald Creek, and Siebert Creek.